

Shailinder Singh FRCS (Paed. Surg.) DM MBA
Consultant Paediatric Surgeon

PS05lite Umbilical Hernia Repair (child)

Expires end of December 2012
Issued January 2012

You can get more information and share your experience at www.aboutmyhealth.org
Tell us how useful you found this document at www.patientfeedback.org



www.rcseng.ac.uk

www.bads.co.uk

www.baps.ac.uk

www.rcsed.ac.uk

This document will give you information about an umbilical hernia repair. If you have any questions, you should ask your GP or other relevant health professional.

What is an umbilical hernia?

An umbilical hernia is caused by a weakness in the layer of muscle of the abdominal wall, just behind the umbilicus (see figure 1). All babies have umbilical hernias when they develop in the womb. Most close in the first three years after birth.



Figure 1

Position of an umbilical hernia

If your child has a hernia, you may notice a swelling, particularly when they cry or strain. Umbilical hernias can be dangerous because the intestines or other structures within the abdomen can get trapped and have their blood supply cut off (strangulated hernia). Although umbilical hernias hardly ever cause these complications in childhood, they are more likely to do so in adulthood.

What are the benefits of surgery?

Your child should no longer have the hernia. Surgery should prevent your child from having any of the serious complications that a hernia can cause in adult life.

Are there any alternatives to surgery?

It is safe to see if the hernia will close without an operation.

If your child is over three years old and they still have an umbilical hernia, the hernia is unlikely to close on its own.

What does the operation involve?

The operation is performed under a general anaesthetic. The operation usually takes between half an hour and three-quarters of an hour.

Your surgeon will make a small cut around half the umbilicus and close the opening in the tough layer of the abdominal wall with strong stitches.

What complications can happen?

1 General complications

- Pain
- Bleeding
- Infection of the surgical site (wound)
- Unsightly scarring

2 Specific complications

- Developing a lump under the wound
- Injury to structures within the hernia
- Unsightly appearance

How soon will my child recover?

Your child should be able to go home the same day.

Your child should be able to return to school after one week but should avoid strenuous activities for six weeks.

Most children make a full recovery. However, the hernia may come back.

Summary

An umbilical hernia is a common condition. If your child is over three years old, surgery is usually recommended to prevent serious complications that can happen in adult life.

Acknowledgements

Author: Mr Shailinder Singh DM FRCS (Paed. Surg.) and Mr Jonathan Sutcliffe FRCS

Illustrations: Medical Illustration Copyright © 2011 Nucleus Medical Art. All rights reserved. www.nucleusinc.com

This document is intended for information purposes only and should not replace advice that your relevant health professional would give you.